

ABSTRACT

MATRIX METALLOPROTEINASE INHIBITORS

5

Compounds are provided that bind allosterically to the catalytic domain of MMP-13 and comprise a hydrophobic group, first and second hydrogen bond acceptors and at least one, and preferably both, of a third hydrogen bond acceptor and a second hydrophobic group. Cartesian coordinates for centroids of the above
10 features are defined in the specification. When the ligand binds to MMP-13, the first, second and third (when present) hydrogen bond acceptors bond respectively with Thr245, Thr 247 and Met 253, the first hydrophobic group locates within the S1' channel of MMP-13 and the second hydrophobic group (when present) is relatively open to solvent. The compounds specifically inhibit the matrix
15 metalloproteinase-13 enzyme and thus are useful for treating diseases resulting from tissue breakdown, such as heart disease, multiple sclerosis, arthritis, atherosclerosis, and osteoporosis.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100